



**SOCIAL DETERMINANTS OF HEALTH FOR
PUBLIC HOUSING RESIDENTS:
DIABETES**

Using data and maps created by National Center for Health in Public Housing (NCHPH) and other national data sources, this publication is one in a series that identifies the prevalence of social factors and population health indicators that affect public housing residents. It is intended for non-clinical health center staff, decision makers, and public housing stakeholders.

SOCIAL DETERMINANTS OF PUBLIC HOUSING RESIDENTS: DIABETES

In 2019, Public Housing Primary Care (PHPC) Health Centers provided comprehensive medical and behavioral health services to 856,191 patients living in or immediately accessible to public housing.¹ The majority of those patients lived in poverty (78%), were uninsured (20%), or were enrolled in some type of public health insurance (54% on Medicaid.)²


Public housing residents often have complex health issues that are impacted by where they live. In general, individuals that receive assistance from the U.S. Department of Housing and Urban Development (HUD) are more likely to have chronic health conditions and are higher utilizers of health care than the rest of the U.S. population, even more so than other low-income renters.

35.8% are in fair or poor health vs. 13.8% of other adults
71% are overweight or obese vs. 64% of other adults
61% have a disability vs. 35.4% of other adults
17.6% have diabetes vs. 9.5% of other adults³

Social, economic, and environmental factors, such as income, access to high quality health care services, and safe and healthy environments play a large role in determining the health status of public housing residents. For example, community assets, such as healthy food outlets and safe places to engage in physical activity, can determine diet and exercise, and in turn, obesity and diabetes rates. The map below shows the location of PHPC Health Centers across the country and the corresponding rates of diabetes in the county where they are located⁴.


60.5% of counties with PHPC Health Centers have high diabetes rates, defined as rates higher than the national average.

Diabetes Prevalence by County



Diabetes is not only more prevalent among public housing residents, it is also more severe. The map below shows the percent of diabetic patients seen at PHPC Health Centers with uncontrolled diabetes, defined as HbA1c levels greater than 9. HbA1c provides a reliable measure of chronic hyperglycemia and correlates well with the risk of long-term diabetes complications. Nationally, about 18% of patients with diabetes have HbA1c levels greater than 9 compared to 32% of diabetic patients seen at PHPC Health Centers⁶. Approximately 90% of PHPC Health Centers have a higher percentage of patients with uncontrolled diabetes compared to the national average.

Uncontrolled Diabetes by County




There are many reasons why diabetes is so pervasive among public housing residents. Limited income, lack of transportation, and access to healthy, affordable foods is a challenge for low income populations. While 15% of the country report they do not have enough food for an active, healthy lifestyle, low income populations are more likely to live in food deserts⁷. A map from the U.S. Department of Agriculture shows that access to food is more challenging for low income populations living in urban and rural areas⁸.

Physical activity is also important for the prevention and management of diabetes and prediabetes. A geospatial analysis by the University of Wisconsin Population Health Institute finds that 87.5% of the population living in counties where PHPC Health Centers are located have access to exercise opportunities through parks and recreational facilities⁹. However, 22% report that they do not engage in any leisure time physical activity¹⁰. There are other factors that can influence decisions to be physically active. For public housing residents, one of those issues is safety.

A meta-analysis by Rees-Punia on the relationship between crime, perceived safety, and physical activity shows that individuals living in areas where they felt unsafe or in areas with higher crime had a 28% reduced odds of achieving higher levels of physical activity¹¹. Rates of violent crime are higher in areas where PHPCs are located compared to the national average. There is an average of 506 violent crimes per 100,000 population in PHPC counties compared to the national average of 386 per 100,000¹².

Violent Crime Rate (per 100k pop) by County



Violence, crime, and the cumulative stress of poverty, unemployment, and other social determinants, impact the mental health of public housing residents. Public housing residents are more likely to report feelings of anxiety, depression, and hopelessness, and those feelings are more likely to interfere with daily activities compared to the general population¹³. Mental health is a critical component to the prevention and management of diabetes because stress informs diet, metabolism, and other health behaviors, such as smoking.

“There is an average of **506** violent crimes per 100,000 population in PHPC counties compared to the national average of 100,000.”

SERIOUS PSYCHOLOGICAL DISTRESS




DISTRESS WITH MENTAL HARDSHIP



Smoking rates have declined over the last decade, but they still remain high for this patient population. About 33.5% of public housing residents smoke compared to 22% of the general population¹⁴. Smoking increases the risk of developing diabetes, makes diabetes harder to control, and increases risk for developing diabetic complications, such as heart and kidney failure, retinopathy, and leg or foot amputations¹⁵.

Adult Smokers by County



Implications

A comprehensive diabetes prevention and management program that addresses the social determinants of health is critical for public housing residents. Health care professionals should consider and address lack of healthy food options, safe places to exercise, community violence, and the stress of living in poverty when developing a plan with their patients.

Many PHPC Health Centers attempt to offset the impact of these determinants through the use of mobile units that provide fresh fruits and vegetables, community gardens and nutrition education programs, exercise and wellness programs with dynamic multilingual staff, flexible hours, transportation, and by collecting and monitoring social determinants of health data for their patient population. For more information on best practices on diabetes prevention and management strategies, please visit www.nchph.org.

SOURCES:

1. Health Resources and Services Administration. (2020) 2019 National Health Center Data: National Public Housing Primary Care Grantee Program Data.
2. Ibid.
3. U.S. Department of Housing and Urban Development. (2017) A Health Picture of HUD-Assisted Adults, 2006-2012. Washington: U.S. Department of Housing and Urban Development.
4. Centers for Disease Control and Prevention. (2013) Diabetes Interactive Atlas. Retrieved from <https://www.cdc.gov/diabetes/atlas/countydata/atlas.html>
5. Health Resources and Services Administration. (2018) 2018 Uniform Data System Public Housing Primary Care Grantee Program Data.
6. National Center for Health Statistics. (2016) Chapter 8: Diabetes. Healthy People 2020 Midcourse Review. Hyattsville: U.S. Department of Health and Human Services; Health Resources and Services Administration. (2016) 2016 National Health Center Data: National Public Housing Primary Care Grantee Program Data. Retrieved from <https://bphc.hrsa.gov/uds/datacenter.aspx?fd=ph>.
7. Feeding America. (2018) Map the Meal Gap: Food Insecurity in America. Retrieved from <http://map.feedingamerica.org/>; U.S. Department of Agriculture, Economic Research Service. (2017) Food Access Research Atlas. Retrieved from <https://www.ers.usda.gov/data-products/food-access-research-atlas/go-to-the-atlas.aspx>
8. U.S. Department of Agriculture, Economic Research Service. (2017) Food Access Research Atlas. Retrieved from <https://www.ers.usda.gov/data-products/food-access-research-atlas/go-to-the-atlas.aspx>
9. Robert Wood Johnson Foundation and the University of Wisconsin Population Health Institute. (2018) County Health Rankings and Roadmaps. Retrieved from <http://www.countyhealthrankings.org/>
10. Centers for Disease Control and Prevention. (2013) Diabetes Interactive Atlas. Retrieved from <https://www.cdc.gov/diabetes/atlas/countydata/atlas.html>
11. Rees-Punia, E., Hathaway, E.D., Gay, J.L. (2018) Crime, perceived safety, and physical activity: A meta-analysis.
12. Preventive Medicine, 111: 307-313.
13. Federal Bureau of Investigation. (2017) Uniform Crime Reporting 2012-2014. Retrieved from <https://ucr.fbi.gov/>
14. U.S. Department of Housing and Urban Development. (2017) A Health Picture of HUD-Assisted Adults, 2006-2012. Washington: U.S. Department of Housing and Urban Development.
15. Ibid.
16. Centers for Disease Control and Prevention. (2018) Tips from Former Smokers: Smoking and Diabetes. Retrieved from <https://www.cdc.gov/tobacco/campaign/tips/diseases/diabetes.html>



SOCIAL DETERMINANTS OF HEALTH FOR
PUBLIC HOUSING RESIDENTS:
DIABETES

Using data and maps created by National Center for Health in Public Housing (NCHPH) and other national data sources, this publication is one in a series that identifies the prevalence of social factors and population health indicators that affect public housing residents. It is intended for non-clinical health center staff, decision makers, and public housing stakeholders.

NCHPH
MAY 2020

DISCLAIMER:

National Center for Health in Public Housing

The mission of National Center for Health in Public Housing (NCHPH) is to strengthen the capacity of federally-funded Public Housing Primary Care (PHPC) health centers and other health center grantees caring for public housing residents by providing training, technical assistance and research. The PHPC program is built on a foundation of collaboration between Health Centers, Public Housing Agencies, and residents. *For more information visit www.nchph.org/interactivemaps.*

This project is supported by the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) as part of an award totaling \$1,824,000 with 0% financed with non-governmental sources. The contents are those of the author(s) and do not necessarily represent the official views of, nor an endorsement, by HRSA, HHS, or the U.S. Government. For more information, please visit HRSA.gov.